

# MAGNETIZATION LOSSES IN HIGH TEMPERATURE SUPERCONDUCTING TAPES DUE TO AC AND DC MAGNETIC FIELDS

Eduard Maievskiy, Marian Cizek\*


Division of Low Temperature and Superconductivity

\*-also at Wrocław University of Technology

# Outline

- 1. Application of superconducting materials**
- 2. Object of studying**
- 3. AC-losses**
- 4. Experimental method and results**

# Application of Superconducting Materials




Eduard Maievskiy  
 Marian Cizek

Institute of Low Temperatures and Structure Research  
 Polish Academy of Science, Wrocław, Poland



# HTSC-tapes

## Cross-section of Sumitomo BSCCO-2223 composite tape type H



Eduard  
Maievskiy




Marian  
Ciszek

Institute of Low Temperatures and Structure Research  
Polish Academy of Science, Wrocław, Poland



# HTSC-tapes

## Cross-section of SCS 4050 2G HTSC tape produced by SuperPower



Eduard  
Maievskiy

Marian  
Ciszek

Institute of Low Temperatures and Structure Research  
Polish Academy of Science, Wrocław, Poland



# Main contributions to AC losses


1. Bulk pinning losses:

$$Q_b \propto \vec{E} \cdot \vec{J}$$



2. Annihilation losses: *vortices of opposite magnetic sense annihilate, one pair liberate energy*

$$Q_a \propto 2H_{c1} \cdot \phi_0$$



3. Surface losses: *arising from a surface barrier*

$$Q_s \propto \Delta H_{eff} \cdot \Phi$$

## Main contributions to AC losses



4. Flux flow losses: *when pinning is very low;*

$$\vec{E}_{ff} \propto \rho_{ff} \cdot \vec{J}; \quad \rho_{ff} \propto \rho_n \cdot \frac{B}{B_{c2}} \quad W_{ff} = f(b_0^2, \omega)$$

5. In HTSC, additionally, contributions from -flux creep, granularity, anisotropy, grain boundaries

6. In composites: *coupling between filaments, eddy currents in matrix*

# Race-Track Coil




Eduard Maievskiy  
Marian Cizek


Institute of Low Temperatures and Structure Research  
Polish Academy of Science, Wrocław, Poland



# Measurement equipment



# Measurement equipment




$$B(x) = \frac{\mu_0 I}{2\pi} \left( \frac{w}{x^2 - \left(\frac{w}{2}\right)^2} \right)$$



$$B(x) = \frac{m}{l} \cdot \left( \frac{1}{2\pi x^2} \right)$$

$$\Phi\left(\frac{p}{2}\right) = -\frac{m}{\pi p} \Rightarrow \Phi(p) = -\frac{2m}{\pi p}$$

$$V(t) = \frac{2}{\pi p} \frac{\partial m}{\partial t}$$



$$m(t) = m' \cos \omega t + m'' \sin \omega t$$

$$m''(t) = \frac{V''(t) \pi p}{2 \omega}$$

$$P = \frac{1}{L} \oint H dm = \frac{1}{L} \int_0^T H(t) \frac{\partial m}{\partial t} dt$$



$$P = \frac{\pi p}{2Lf} H_{rms} V''_{rms}$$

Loop has to be extended from the edge of a tape


M Iwakuma,... *SUST* **16** (2003) 545  
 D N Nguyen,...*SUST* **19** (2006) 1010  
 Y Yang,...*J. App. Phys.* **96** (2004) 2141

Eduard Maievskyi  
 Marian Ciszek

Institute of Low Temperatures and Structure Research  
 Polish Academy of Science, Wroclaw, Poland



# Energy dissipation in BSCCO-2223 tape AC-field only





Eduard Maievskiy  
Marian Cizek

Institute of Low Temperatures and Structure Research  
Polish Academy of Science, Wrocław, Poland



# Energy dissipation in YBCO-123 tape (AC-field only)





Eduard Maievskiy  
Marian Cizek

Institute of Low Temperatures and Structure Research  
Polish Academy of Science, Wrocław, Poland



# Energy dissipation in YBCO-123 tape (AC-field only)

## Loss model for a thin film (perpendicular field)




$$J_y = -\left(\frac{\partial H_z}{\partial x} - \frac{\partial H_x}{\partial z}\right) \quad \text{when } d \ll a$$

$$J_y(x) = -\frac{c}{4\pi d} [B_x(x, z = d/2) - B_x(x, z = -d/2)]$$

$$Q_m = \frac{4\pi}{\mu_0} a^2 b_0^2 \cdot g(x)$$

$$g(x) = \frac{1}{x} \left[ \frac{2}{x} \ln \cosh(x) - \tanh(x) \right]$$



$$Q_m \propto a^2 b_0^4$$

$$Q_m \propto I_c a^2 b_0$$

$$x = b_0 / B_c$$

$$B_c = \mu_0 I_c / 2\pi a$$



E.H. Brandt and M. Indenbom, PRB 48 (1993) 12893  
J.R. Clem, E. Zeldov, PRB 49 (1994) 9802.

Eduard Maievskyi  
Marian Cizek

Institute of Low Temperatures and Structure Research  
Polish Academy of Science, Wrocław, Poland



# Energy dissipation in YBCO-123 tape (AC-field only)







Eduard Maievskiy  
Marian Cizek

Institute of Low Temperatures and Structure Research  
Polish Academy of Science, Wrocław, Poland



# Hysteretic loss minimum in BSCCO-2223 tape







Eduard Maievskyi  
Marian Cizek

Institute of Low Temperatures and Structure Research  
Polish Academy of Science, Wrocław, Poland



# Hysteretic loss minimum in YBCO-123 tape




Eduard Maievskiy  
Marian Cizek

Institute of Low Temperatures and Structure Research  
Polish Academy of Science, Wrocław, Poland



## Clem model



$W'_L$  - total losses  
 $W'_S$  - surface losses  
 $W'_{bp}$  - bulk pinning losses  
 $W'_{ba}$  - bulk annihilation losses

John R. Clem, „Theory of AC losses in type II superconductors with a field-dependent surface barrier”,  
J. Appl. Phys. 50(5), May 1979

*Thank you for attention!*

Eduard  
Maievskiy

Marian  
Ciszek

Institute of Low Temperatures and Structure Research  
Polish Academy of Science, Wrocław, Poland

